Supplementary Table S1. Distribution of the histologic components within each histologic group

	tub	por	pcc	sig
Early gastric cancer (n = 459)				
tub ≥10% (n = 398)	100.0 (89.7–100.0)	0.0 (0.0–8.5)	0.0 (0.0–0.0)	0.0 (0.0–0.0)
por ≥10% (n = 120)	46.0 (16.5–75.1)	30.0 (19.5–55.4)	0.0 (0.0–0.1)	0.0 (0.0–5.0)
pcc ≥10% (n = 43)	0.0 (0.0–6.4)	3.2 (0.0–21.5)	37.0 (24.5–58.5)	32.4 (4.2–62.7)
sig ≥10% (n = 66)	0.0 (0.0–5.0)	5.8 (0.0–23.3)	6.1 (0.0–33.7)	60.3 (34.2–90.0)
Advanced gastric cancer (n = 141)				
tub ≥10% (n = 79)	60.4 (32.6–88.8)	28.7 (6.2–41.7)	0.0 (0.0–5.0)	0.0 (0.0–0.0)
por ≥10% (n = 81)	24.3 (5.0–55.5)	39.6 (28.7–70.0)	0.0 (0.0–26.7)	0.0 (0.0–1.8)
pcc ≥10% (n = 67)	0.2 (0.0–19.5)	7.0 (0.3–35.6)	65.4 (35.9–89.2)	3.9 (0.0–11.7)
sig ≥10% (n = 25)	0.0 (0.0–0.3)	4.4 (0.6–27.2)	50.8 (10.0–68.7)	28.9 (21.5–49.2)
Total cohort (n = 600)				
tub ≥10% (n = 477)	100.0 (75.4–100.0)	0.0 (0.0–18.0)	0.0 (0.0–0.0)	0.0 (0.0–0.0)
por ≥10% (n = 201)	39.9 (10.2–70.3)	36.5 (21.1–60.0)	0.0 (0.0–5.9)	0.0 (0.0–2.9)
pcc ≥10% (n = 110)	0.0 (0.0–12.0)	4.7 (0.0–23.9)	50.3 (30.1–78.7)	6.5 (0.8–35.4)
sig ≥10% (n = 91)	0.0 (0.0–4.6)	5.0 (0.0–25.3)	10.0 (0.0–45.8)	49.8 (28.5–77.0)

Values are median (interquartile range) percentage of each component within the whole tumor.

EGC, early gastric cancer; tub, tubular adenocarcinoma, well to moderately differentiated; por, tubular adenocarcinoma, poorly differentiated; pcc, poorly cohesive carcinoma, not otherwise specified; sig, signet-ring cell carcinoma.